



This is a repository copy of *ApoE ϵ 4 allele related alterations in hippocampal connectivity in early Alzheimer's disease support memory performance.*

White Rose Research Online URL for this paper:
<http://eprints.whiterose.ac.uk/112432/>

Version: Accepted Version

Article:

De Marco, M., Vallelunga, A., Meneghello, F. et al. (4 more authors) (2017) ApoE ϵ 4 allele related alterations in hippocampal connectivity in early Alzheimer's disease support memory performance. Current Alzheimer Research . ISSN 1567-2050

<https://doi.org/10.2174/1567205014666170206113528>

Reuse

Unless indicated otherwise, fulltext items are protected by copyright with all rights reserved. The copyright exception in section 29 of the Copyright, Designs and Patents Act 1988 allows the making of a single copy solely for the purpose of non-commercial research or private study within the limits of fair dealing. The publisher or other rights-holder may allow further reproduction and re-use of this version - refer to the White Rose Research Online record for this item. Where records identify the publisher as the copyright holder, users can verify any specific terms of use on the publisher's website.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.



eprints@whiterose.ac.uk
<https://eprints.whiterose.ac.uk/>

ApoE ε4 allele related alterations in hippocampal connectivity in early Alzheimer's disease support memory performance

Matteo De Marco^{a,b}, Annamaria Vallelunga^c, Francesca Meneghello^b, Susheel Varma^d, Alejandro Frangi^{d,e,f}, Annalena Venneri^{a,b}, for the Alzheimer's Disease Neuroimaging Initiative*

^a University of Sheffield, Sheffield Teaching Hospital NHS Foundation Trust, Medical School, Beech Hill Road, S10 2RX, Sheffield, UK;

^b IRCCS Fondazione Ospedale San Camillo, via Alberoni 70, 30126, Lido, Venice, Italy;

^c Neuroscience Section, Department of Medicine and Surgery, Center for Neurodegenerative Diseases (CEMAND), University of Salerno, Salerno, Italy;

^d Centre for Computational Imaging & Simulation Technologies in Biomedicine (CISTIB), Department of Electronic and Electrical Engineering, University of Sheffield, Sheffield, UK

^e Insigneo Institute for in silico Medicine, University of Sheffield, Sheffield, UK

^f Department of Electronic and Electrical Engineering, University of Sheffield, Sheffield, UK

*Data used in preparation of this article were obtained from the Alzheimer's Disease Neuroimaging Initiative (ADNI) database (adni.loni.usc.edu). As such, the investigators within the ADNI contributed to the design and implementation of ADNI and/or provided data but did not participate in analysis or writing of this report.

A complete listing of ADNI investigators can be found at: http://adni.loni.usc.edu/wp-content/uploads/how_to_apply/ADNI_Acknowledgement_List.pdf

Address for correspondence:

Prof. Annalena Venneri

Department of Neuroscience - Medical School, University of Sheffield

Beech Hill Road, Royal Hallamshire Hospital, N floor, room N130

Sheffield - S10 2RX - United Kingdom

Acknowledgements

The BioBank of Rare and Neurorehabilitative Diseases, IRCCS Fondazione Ospedale San Camillo provided us with specimens. The authors thank Cristina Pilosio, Jessica Rigon, Francesca Bevilacqua, Davide Duzzi, Elisabetta Tasca, and all personnel in the MRI unit of the IRCCS Fondazione Ospedale San Camillo in Venice, Italy for their various contribution to patient assessment, imaging acquisitions and planning of image analyses. The authors also thank Leandro Beltrachini and Ernesto Coto for their help with ADNI image processing and modelling. As part of the STEPS procedures of hippocampal segmentation, the GPU card used for this research was donated by the NVIDIA Corporation. This study has been supported by grant No. 42/RF-2010-2321718 from the Italian Ministry of Health to AV. MDM and SV are employed through funding from the European Union Seventh Framework Programme (FP7/2007 – 2013) under grant agreement no. 601055, VPH-DARE@IT to AV and AF.

For the purpose of post-hoc analyses, data collection and sharing for this project was funded by the Alzheimer's Disease Neuroimaging Initiative (ADNI) (National Institutes of Health Grant U01 AG024904) and DOD ADNI (Department of Defense award number W81XWH-12-2-0012). ADNI is funded by the National Institute on Aging, the National Institute of Biomedical Imaging and Bioengineering, and through generous contributions from the following: AbbVie, Alzheimer's Association; Alzheimer's Drug Discovery Foundation; Araclon Biotech; BioClinica, Inc.; Biogen; Bristol-Myers Squibb Company; CereSpir, Inc.; Cogstate; Eisai Inc.; Elan Pharmaceuticals, Inc.; Eli Lilly and Company; EuroImmun; F. Hoffmann-La Roche Ltd and its affiliated company Genentech, Inc.; Fujirebio; GE Healthcare; IXICO Ltd.; Janssen Alzheimer Immunotherapy Research & Development, LLC.; Johnson & Johnson Pharmaceutical Research & Development LLC.; Lumosity; Lundbeck; Merck & Co., Inc.; Meso Scale Diagnostics, LLC.; NeuroRx Research; Neurotrack Technologies; Novartis Pharmaceuticals Corporation; Pfizer Inc.; Piramal Imaging;

Servier; Takeda Pharmaceutical Company; and Transition Therapeutics. The Canadian Institutes of Health Research is providing funds to support ADNI clinical sites in Canada. Private sector contributions are facilitated by the Foundation for the National Institutes of Health (www.fnih.org). The grantee organization is the Northern California Institute for Research and Education, and the study is coordinated by the Alzheimer's Therapeutic Research Institute at the University of Southern California. ADNI data are disseminated by the Laboratory for Neuro Imaging at the University of Southern California.

Abstract

Background. Whether the presence of the Apolipoprotein E ϵ_4 allele modulates hippocampal connectivity networks in abnormal ageing has yet to be fully clarified.

Objective. Allele-dependent differences in this pattern of functional connectivity were investigated in patients with very mild neurodegeneration of the Alzheimer's type, carriers and non-carriers of the ϵ_4 allele.

Method. A seed-based connectivity approach was used. The two groups were similar in demographics, volumetric measures of brain-structure, and cognitive profiles.

Results. ϵ_4 carriers had increased connectivity between the seed area in the left hippocampus and 1) a left insular/lateral prefrontal region and 2) the contralateral right parietal cortex. Moreover, hippocampus-to-parietal connectivity in the group of ϵ_4 carriers was positively associated with memory performance, indicating that the between-group difference reflects compensatory processes. Retrospective analyses of functional connectivity based on patients from the ADNI initiative confirmed this pattern.

Conclusion. We suggest that increased connectivity with extra-DMN areas reflects both compensatory recruitment of additional areas, and pathological intertwining between the DMN and the salience network as part of a global ϵ_4 -dependent circuitual disruption. These differences indicate that the ϵ_4 allele is associated with a more profound degree of DMN network breakdown even in the prodromal stage of neurodegeneration.

Keywords

Functional connectivity; Disinhibition hypothesis; Posterior cingulate; Hippocampus; Alzheimer's disease; Network disruption;

Running Title

Hippocampal connectivity in ApoE ϵ_4 -positive early-AD

1. Introduction

The brain default-mode network (DMN) is a functional resting-state circuit that normally activates while a person does not engage in any explicit mental task [1]. It includes midline kernels localised in the posterior cingulate and medial prefrontal cortices, which are connected functionally with the inferior parietal lobule, the lateral temporal cortex, and the hippocampal formation [2]. Alzheimer's disease (AD) causes a global disruption of functional connectivity within the DMN [3]. Further segmentation of the circuit into sub-components suggests that AD down-regulates connectivity within the structures located in the posterior portion of the DMN and up-regulates connectivity of the prefrontal hubs [4-5]. This process begins in prodromal AD, when the disease is characterised by a transitional phase of Mild Cognitive Impairment (MCI) [6-7]. Within the set of regions included within the DMN, the hippocampus plays a distinguishing role. In fact, hippocampal subfields are subjected to a well-established volumetric loss along the timeline of AD [8], and this shrinkage is predictive of conversion from MCI to the dementia stage [9]. Moreover, when compared with healthy controls, patients diagnosed with amnesic MCI (thus suggestive of potential AD) show disrupted connectivity, with pathological up-regulation of connectivity within the hippocampal formation and between the hippocampus and the posterior-cingulate/precuneus region [10]. Modifications of memory-associated patterns of hippocampal activation have been also reported in MCI [11], and this aspect was found to be associated with cortical thinning of "signature regions" of AD including frontal, temporal and parietal cortices [12].

The ϵ_4 isoform of the Apolipoprotein E (ApoE) gene is a well-established risk factor for the development of the sporadic late-onset forms of AD [13-14], and is associated with a younger age of disease onset [15]. The homonymous peptide coded by the gene plays a crucial role in lipoprotein metabolism and neurobiology [16]. As for the latter set of functions, the ApoE ϵ_4 allele appears responsible for a large number of detrimental effects on neuronal and synaptic function, in comparison with the "standard" ϵ_3 isoform [17], and appears to exert its impact on cell biology both either in the presence or in the absence of neurodegeneration [18]. Following the Imaging Genetics model, the direct effect of the various ApoE isoforms on cellular mechanisms translates into indirect, yet consequent, effects of the genotype on the development of brain structure and brain function [19]. Within this latter category of variables, a large number of studies on healthy adults have found that the ϵ_4 isoform is associated with alterations of the DMN both in healthy ageing and young adulthood [20-28], in correspondence or even prior to the initiation of the neuropathological cascade seen in AD. Conversely, the study of the impact of the ϵ_4 allele on functional connectivity in the symptomatic stages of AD has been scarce. A few studies investigated electro- and magneto-

encephalographic connectivity in samples of MCI and mildly-to-moderately demented AD patients. These revealed that the ϵ_4 isoform is associated with decreased levels of functional connectivity as measured by various proxies of connectivity such as signal coherence, synchronisation likelihood, and ROI-based lagged-phase synchronisation [29-32]. Aside from well-established limits in spatial resolution, however, these techniques do not allow a specific focus on all major DMN hubs affected by AD, as these are located in regions that are not easily capturable by measurements obtained at the scalp level. Overcoming these methodological limitations, two very recent rest-fMRI studies found that the DMN of ϵ_4 patients diagnosed with early-stage AD is significantly down-regulated [33-34]. Despite the limited number of studies, this body of evidence indicates that the presence of the ϵ_4 allele impacts negatively on DMN integrity even after the onset of a clinically-established symptomatology of dementia. Nevertheless, it is still undetermined whether variability for the ApoE genotype is associated with a distinctive signature of disruption of hippocampal functional connectivity in the prodromal phases of the disease, when the person still retains their daily life independence. The hippocampus, harshly affected in AD, is of particular interest in this early phase of the disease. In fact, this stage is crucial because it represents the earliest moment of subjective/objective awareness of the presence of a possible neurodegenerative disease. In addition, published studies suggest that this clinical stage is associated with a high degree of retained mechanisms of neuroplasticity, sufficient to induce remarkable changes in brain functioning [e.g. 35-36]. It is thus of paramount importance to characterise and interpret appropriately allele-dependent differences in the connectivity of a core region like the hippocampus during such a clinically relevant stage, as qualitative differences in the functional architecture of the brain might translate into qualitative differences in the efficacy of treatments between carriers and non-carriers (e.g. a memory-enhancing training).

In this study we investigated the fMRI network signature of the ApoE ϵ_4 allele in hippocampal connectivity among patients suffering from very mild AD. To do so, we implemented seed-based connectivity methods in a sample of patients ϵ_4 carriers and in a sample of ϵ_4 non-carriers. We hypothesised that allele-dependent differences would exist between the two groups, and that the presence of the ϵ_4 allele would be associated with a signature of connectivity involving associative areas which sustain high-order cognitive processing.

2. Material and methods

2.1. Participants

Sixty-one patients were referred to neurological examination between 2011 and 2014 because of suspected incipient cognitive decline. On that occasion, all patients agreed on completing cognitive assessment and an MRI procedures (detailed below). A proportion of these patients was diagnosed with very mild dementia of the AD type [37], while a proportion received a diagnosis of MCI [38]. This latter group was followed-up over time, and progression of disease and conversion were monitored until early 2015. Based on this longitudinal neurological monitoring, only patients who showed evidence of a clinically-established progression towards AD dementia at follow up were included in this study.

A neurological examination served to rule out the presence of major exclusion criteria, which were set as follows: a significant disease at clinical level, history of transient ischemic attacks, a diagnosis of vascular brain disease of clinical severity (e.g. the presence of chronic cerebrovascular disease as main aetiology), a structural MRI revealing a different diagnostic entity which could otherwise explain the presence of cognitive symptoms, presence/diagnosis of uncontrolled seizures, peptic ulcer, cardiovascular disease, sick sinus syndrome, neuropathy with conduction difficulties, significant disabilities, proof of abnormal baseline levels of folates, vitamin B12 or thyroid-stimulating hormone, a significant psychiatric condition, consumption of drugs for research purposes or with toxic effects to internal organs.

After genetic assays, all AD patients carrying at least one copy of the ϵ_4 allele ($n = 15$) were enrolled. None of these had a $\epsilon_4\epsilon_4$ or a $\epsilon_2\epsilon_4$ genotype. A group of non-carriers ($\epsilon_3\epsilon_3$ only) was then selected from the pool of remaining patients to match the two groups as closely as possible for demographic characteristics. Other non ϵ_4 genotypes were not included in the control sample to avoid contamination of protective factors such as in the case of the ϵ_2 genotype, and in an attempt to minimise variance in the control patient group. All participants were Caucasian and were inhabitants of one of the islands within the Venetian lagoon. This study was carried out according to the Declaration of Helsinki and was approved by the Institutional Review Board of the IRCCS Fondazione Ospedale San Camillo (Venice, Italy). Written informed consent was obtained from each study participant.

An extensive battery of neuropsychological tests was administered to each patient as part of the initial diagnostic classification procedures. This included tests assessing short- and long-term verbal and non-verbal memory, attention, naming by confrontation, logical abstract reasoning, verbal fluency and visuoconstructional abilities (see Table 1 for details).

2.2. MRI Acquisition, Preprocessing and Analysis

A structural 3D T1-weighted brain scan and two resting-state fMRI runs were acquired on a 1.5 T Philips Achieva system, and preprocessed and analysed using Statistical Parametric Mapping (SPM) 8 software (Wellcome Trust Centre for Neuroimaging, London, UK) running in Matlab R2011b (Mathworks Inc., UK). T2-weighted and FLAIR-weighted sequences were also included in the protocol to verify neuro-anatomical exclusion criteria and suitability for inclusion in the study. A senior neuroradiologist reviewed each anatomical scan to ascertain study compatibility. Participants were asked to remain as still as possible for the full duration of the scan. No stimuli were presented.

Preprocessing of T1-weighted images was carried out using a standard Voxel-Based Morphometry approach [39]. Native-space volumes of grey matter, white matter, and cerebrospinal fluid were obtained to calculate individual brain parenchymal volume, total intracranial volume, and tissue-class ratios (grey-matter, white-matter, and brain parenchymal fraction). Modulated and normalised tissue-class maps were then smoothed with a 8 mm full-width at half maximum gaussian kernel. Additionally, native-space T1-weighted images were also segmented to extract the hippocampal maps for further sample characterisation. For this purpose, the STEPS algorithm was implemented [40]. Briefly, this methodology allows an automatic and precise segmentation of the hippocampus by registering each scan to the most appropriate image among a series of available templates. Absolute and ratio-based volumetric properties of left and right hippocampus were thus extracted.

Resting-state fMRI acquisitions were preceded by 20-seconds of dummy scans to allow the scanner to reach a state of electro-magnetic equilibrium. Each run included 120 volumes of T2* weighted echo planar images (repetition time = 2 s, echo delay time = 50 ms, flip angle 90°, voxel dimensions $3.28 \times 3.28 \times 6.00 \text{ mm}^3$, field of view 230 mm). Each volume included 20 contiguous axial slices, acquired in ascending order. Slice-timing was carried out first. Volumes in each run were then realigned and resliced independently. Concurrently, linear and rotational parameters of head motion were estimated by the use of 4th Degree B-Spline interpolation. Volumes were subsequently

normalised using the first realigned volume as source image to match the SPM 8 echoplanar template, and voxel size was re-dimensioned to $2.0 \times 2.0 \times 2.0 \text{ mm}^3$. Images were then band-pass filtered at 0.008 – 0.1 Hz using the REST toolbox [41]. Finally, volume smoothing was carried out with a $6.0 \times 6.0 \times 6.0 \text{ mm}^3$ full-width at half maximum isotropic gaussian kernel.

Hippocampal connectivity was computed by means of seed-based first-level models. Seed regions were devised based on the IBA16 atlas implemented in the WFU-Pickatlas toolbox [42]. Both left and right seeds were loaded on a structural template to ascertain the absence of major spatial misplacements over the temporal horn of the ventricles. By doing so, miscalculations of average seed signal due to the presence of cerebrospinal liquid were minimised. Signal extraction from the two seed regions was carried out using the MarsBaR toolbox [43]. Two additional vectors were extracted from the map of white matter and from that of cerebrospinal fluid. First-level analyses were carried out to obtain individual maps of seed-based connectivity, regressing out the signal from white matter and cerebrospinal fluid, and controlling for in-scanner motion vectors. For inferential analyses, a p value of 0.01 (uncorrected) was set. Age, education levels, MMSE scores, and grey-matter fraction were used as covariates. MMSE scores were included in the model to account for variability of disease severity, as the mechanisms of AD affect the connectivity of the hippocampus [44-45], while grey-matter fraction served as proxy of brain reserve. Of all the output clusters only peaks surviving Family-Wise Error (FWE)-correction at a cluster level were reported as significant to minimise chances for Type I Errors. Peak coordinates were converted into Talairach stereotaxic space thanks to a non-linear transform (<http://imaging.mrc-cbu.cam.ac.uk/downloads/MNI2tal/mni2tal-m>) and interpreted using the Talairach Daemon client (www.talairach.org/client.htm), single-point coordinate search [46-47].

3. Results

There was no significant difference in age, level of education and male/female proportion between the two groups. Group comparisons revealed also no differences in absolute/proportional properties of brain structure. Although global difference in white-matter ratio survived correction for multiple statistical comparisons (with 483 patients having a significantly higher ratio of white matter), the voxel-based analyses revealed no between-group differences neither in the regional maps of grey matter nor in those of white matter. Moreover, all p values indicated that in our sample the two hippocampal regions were comparable between the two groups, as both absolute and relative

(fractional) volumetric values did not differ between ϵ_4 carriers and non-carriers. No between-group difference was found in any of the raw scores obtained over the set of neuropsychological tests, not even when age and education were included in the analyses. No significant differences were found also when age- and/or education corrected scores based on published norms were compared between the two groups. Both groups of patients had had an amnesic onset as established by their neuropsychological profile. Verbal declarative memory was, in fact, the cognitive domain in which both carriers and non-carriers showed performance levels below cut-off. All these between-group comparisons are reported in Table 1.

- Insert Table 1 about here -

Hippocampal connectivity findings are reported in Table 2 and illustrated in **Fig.1**. ϵ_4 carriers showed enhanced connectivity between the left hippocampus and two clusters, one located in the left insula extending to the inferior frontal cortex, and one located in the right inferior parietal lobule, with a peripheral peak located in somatosensory areas. No differences were found in the functional connectivity of the right hippocampus.

- Insert **Fig.1** and Table 2 about here –

In order to clarify whether this allele-dependent difference in the pattern of functional connectivity between the hippocampus and the inferior parietal lobule was beneficial or not, this latter cluster was binarised, and signal extraction was carried out from this region as originally carried out for the two seeds. A seed-to-target index of connectivity was then computed. A coefficient of partial correlation was calculated between seed and target vectors, controlling for the same regressors as with the voxel-based analyses, and a *Fischer's r to z* transformation was then applied. Since the recruited sample was prevalently characterised by amnesic problems, a composite index of memory performance was computed by transforming the corrected scores of four tests investigating various aspects of verbal and visuospatial memory (Rey Complex Figure Test – Recall, Visual Supraspan Test, Prose Memory Test - Global Recall, and Paired Associates Test) into z-scores, which were then averaged for each patient. This variable distributed normally (Shapiro-Wilk test of

10

normality $p = 0.920$). A linear-regression model was then designed for both groups. To test the hypothesis by which this pathway of connectivity predicted memory performance, two blocks were created. The Mini-Mental State Examination score and the fractional volume of the left hippocampus were inputted in the first block to control for potential cognitive and neurostructural confounds, and the z-index of connectivity was included in the second block. In the group of non-carriers, the slope associated with the predictor was not significant ($b = 0.131$; r^2 -change statistic = 0.001; $p = 0.905$), whereas a significant slope was found in the group of carriers ($b = 0.759$; r^2 -change statistic = 0.211; $p = 0.029$; **Fig.2**). This allele-dependent association did not generalise to other cognitive functions. In fact, the strength of this specific pathway of connectivity did not predict executive performance (average of z-transformed corrected scores in the Stroop Test – Time and the Letter Fluency Test) in neither of the two groups.

- Insert **Fig.2** about here -

3.1 Validation of the results in the ADNI cohort

To verify this pattern of group difference, an additional sample of 65 MCI patients was identified (30 with an $\epsilon_4\epsilon_3$ or $\epsilon_4\epsilon_4$ genotype, and 35 with an $\epsilon_3\epsilon_3$ genotype) (see Table 3 for details). These were obtained from the Alzheimer's Disease Neuroimaging Initiative (ADNI) database (adni.loni.usc.edu). The ADNI was launched in 2003 as a public-private partnership, led by Principal Investigator Michael W. Weiner, MD. The primary goal of ADNI has been to test whether serial magnetic resonance imaging (MRI), positron emission tomography (PET), other biological markers, and clinical and neuropsychological assessment can be combined to measure the progression of mild cognitive impairment (MCI) and early Alzheimer's disease (AD). ROI-based analyses were carried out to compute the pattern of functional connectivity between the hippocampus and Brodmann areas. Univariate ANOVAs were then run between the two groups controlling for age, education levels, MMSE and ventricle size. Increased functional connectivity was found in the group of ϵ_4 carriers in the supramarginal gyrus ($p < 0.029$) and in the prefrontal cortex ($p < 0.039$), paralleling our original findings.

- Insert Table 3 about here -

4. Discussion

In this study, differences within the maps of hippocampal connectivity were investigated between a group of early stage AD patients carrying a copy of the ApoE ϵ_4 allele, and a group of patients with a $\epsilon_3\epsilon_3$ genotype. The sample was extensively tested with statistical procedures in order to maximise comparability along the most relevant axes of demographic, neurostructural and cognitive variability.

The findings show that carriers have increased connectivity between the left hippocampus and two clusters, the first of which was centred in the right inferior parietal lobule. Being the hippocampus and the inferior parietal lobule both hubs of the DMN [2], this piece of evidence suggests that the presence of the ϵ_4 allele might be associated with a more preserved DMN in prodromal AD. This trend, however, goes in the opposite direction as that emerged from the studies characterising the DMN when AD is more severe, in which ϵ_4 carriers were instead found to show reduced connectivity within aspects of the DMN [34], and, specifically, between the hippocampus and prefrontal, parietal, and temporal regions [33]. As a consequence, we suggest that the ϵ_4 allele might influence the patterns of hippocampal connectivity with a quadratic tendency. This trend would not come as a novelty. In fact, it follows the longitudinal progression of hippocampal function as reported by a study of task fMRI, in which activation of this region is increased in MCI patients during memory processes, but reduced in AD dementia [11]. Furthermore, recent evidence indicates that there is an inverse association between hippocampal function (as measured by FDG PET metabolism) and DMN connectivity (as estimated by hippocampus-to-precuneus BOLD-signal correlation) in patients with AD dementia, but such association is not visible in MCI [48]. These pieces of evidence indicate that the progressive disruption of DMN and hippocampal connectivity are not linear along the timeline of AD progression. We hereby suggest that the enhanced hippocampus-parietal connectivity seen in the group of ϵ_4 carriers reflects an intensification of this “naturally-occurring” phenomenon. Up-regulation of connectivity between hippocampus and left lateral prefrontal cortex was seen in the group of ϵ_4 carriers. The involvement of the frontal lobe as ApoE-dependent between-group difference has already been reported in studies of healthy individuals, in which augmented connectivity was observed in ϵ_4 carriers between hubs of the DMN and extra areas that normally are not part of this circuit [e.g. 22, 24]. This was accounted for by the

hypothesis of compensatory mechanisms taking over from the AD-dependent disruption of “standard” patterns of connectivity. The idea that a functional reorganisation of regional connectivity occurs in healthy adults with a risk factor for AD is also supported by studies of task-associated fMRI, in which evidence of computational differences has been repeatedly reported, albeit with no constant pattern [49]. Our results suggest that compensatory mechanisms may be triggered in ϵ_4 carriers even after the possible onset of neurodegeneration, or it might be the outcome of the brain over-time coping with the subtle negative effects of this genetic risk factor.

At a first glance, the up-regulated connectivity between the hippocampus and the parietal lobe might also be compensatory in nature, as it consists of an intensification of a pattern of connectivity which is normally visible by default. Despite this straight-forward interpretative remark, there is another potential explanation that needs to be taken into account, which suggests that the evidence of “hippocampal hyperconnectivity” is not necessarily index of compensation. A recent randomised trial found that amnesic MCI patients receiving mild antiepileptic medication showed a significant improvement in memory performance, which associated with significantly reduced hippocampal activation. As a consequence the enhanced magnitude of hippocampal function seen at baseline was interpreted by the authors as a dysfunctional trait [50]. Based on this same interpretational paradigm (albeit transposed, in a speculative way, to the construct of connectivity), it might be suspected that the increased connectivity seen in the DMN of ϵ_4 carriers might be maladaptive in nature, despite being located in a functional pathway where, at least in healthy adults, “more would be better”. Along this plausible line, a study found increased task-based connectivity of hippocampal seeds in amnesic MCI patients compared with healthy adults in a set of regions including the prefrontal, temporal, parietal and limbic lobe [51]. Additionally, many of these pathways of connectivity were inversely associated with cognitive performance, as an indication of the dysfunctional nature of this excessive connectivity [51]. To test the conflicting hypotheses of compensation vs. maladaptive rewiring, we carried out post-hoc analyses to explore the association between the strength of the functional connectivity between the hippocampus and the inferior parietal lobule, and an index of memory performance. While no association was found in the group of non-carriers, a positive association was found in the group of carriers, indicating that, compatibly with a compensatory mechanism, the more connected the two regions, the more productive the memory processing.

There is evidence that increases in functional connectivity seen in AD neurodegeneration might be the result of a pathological interconnection of distinct network patterns. This was reported by Wang and colleagues [34], who, using a graph-theory approach, found decreased levels of inter-network

connectivity among AD patients carriers of the ϵ_4 isoform. Conversely, Seeley and colleagues [52] came up with the same interpretational avenue to explain the spreading (and, therefore, increase) of functional connectivity of the DMN to extra-DMN structures. Neural cerebral cortex circuitry is a biological system characterised by balanced involvement of parallel networks. The DMN is negatively correlated with the salience network, a resting-state circuit whose activation is associated with the integration of sensory processing and internal autonomic-visceral processing [53]. This anticorrelation would reflect an inter-network balance based on mutual inhibition between the two circuits. When the DMN is damaged by AD, this harmonic equilibrium would collapse, and, as a consequence, the anticorrelated network would no longer be inhibited. This disinhibition would trigger enhanced connectivity within the anticorrelated network and reorganisation of connectivity patterns to induce between-network interactions [52]. Published evidence provides additional support for this hypothesis as a potential signature of the ϵ_4 allele. In a study of healthy older individuals, ϵ_4 carriers showed more connectivity within the salience network than non-carriers [21]. Another study of network differences associated with the presence of the ϵ_4 isoform in middle-aged adults found diminished connectivity within structures of the DMN in carriers, including the left hippocampal/parahippocampal complex, the left anterior temporal pole and the dorsomedial prefrontal cortex bilaterally. At the same time decreased anticorrelated connectivity was observed between the posterior cingulate and regions in the salience network. Complementarily, seed-based connectivity devised to estimate the SN revealed increased connectivity with DMN areas in the same group [54]. Consistently with the disinhibition hypothesis, we interpret the enhanced functional connectivity seen in ϵ_4 carriers between the seed in the left hippocampus (part of the DMN) and the left insula, that is one of the main hubs of the salience network, as maladaptive.

These findings support the idea that allele-dependent diversity in resting-state circuitry is detectable even after conversion from healthy ageing to the first symptomatic phases of AD. The various interpretations of differences found between the two groups converge in indicating that patients carriers of the ApoE ϵ_4 allele have additional circuitual damage in comparison with non-carrier individuals. This additional network disruption is supported by the necessity to rely on a higher magnitude of compensation and by the dysfunctional nature of some of the circuitual rewiring. It is noteworthy to highlight that no cognitive differences and no neurostructural discrepancies existed between the two groups of patients. Nonetheless, circuitual breakdown was more profound in the group of ϵ_4 carriers. This may have important implications in the therapeutical management of these patients. Two individuals with comparable cognitive phenotype might have a diverse degree

of hidden circuitual deterioration/dysfunction that is genotype-dependent. A more profoundly damaged neural architecture might be associated to less capacity for neuroplastic changes, and some forms of therapeutic intervention might not be as beneficial as expected.

These results support the view that, despite its non-linear tendency (increase of DMN connectivity seen in the prodromal phases, and decrease of DMN connectivity documented in the later dementia phases), the ϵ_4 -associated network disruption might be independent of disease stage. Recent findings suggest that the neural representations of at least some aspects of certain cognitive functions are subjected to a comprehensive re-organisation in association with the development of MCI [55-56]. This insidious functional “re-moulding” could potentially nullify or at least minimise the detrimental impact of the ϵ_4 allele, which would be superseded by the impact of pathology. Our study concludes that this does not occur, as the impact of genetic variability for the ApoE gene keeps expressing in the form of more profound condition of network disruption even after the onset of prodromal pathological processes. Other findings are in line with this conclusion. There is a strong body of evidence suggesting that ϵ_4 carriers with MCI tend to have smaller hippocampi and amygdalae than patients not carrying the ϵ_4 allele but having similar demographic characteristics [57-60]. Other studies have instead reported volumetric loss in ϵ_4 carriers extending to other cortical and subcortical areas [61-64], although some have suggested that non-carriers can cope with a much more pronounced and extensive brain volume loss before manifesting the same level of cognitive disruption [64]. In addition, there is evidence that MCI patients carrying the ϵ_4 isoform have reduced cortical metabolism bilaterally in the precuneus, the superior temporal gyrus and the inferior parietal lobule [65]. Albeit being in line with our findings, all these studies were based on a localisation-based approach. This type of approach is not as suitable as a connectivity-based framework to characterise cellular and synaptic disruption, and its association to the underlying cognitive functions [66]. Moreover, it is worth noting that evidence emerging from connectivity analyses should not be taken for granted as the natural consequence of the aforementioned localisation-based differences reported in the literature. Indeed, the parallel study of the impact of the ϵ_4 allele on structural connectivity of AD patients (which describes different, yet theoretically contiguous aspects of signal propagation) has led to incongruous findings [67-68].

Although a “classical” view indicates that the left hippocampus would be mainly involved in verbal memory processes (as opposed to the right hippocampus, involved instead mainly in visuospatial memory processes), recent evidence found instead no evidence of such lateralisation [69]. On this note, the creation of a memory composite score based on both verbal and visuospatial memory

performance would be a better modality-independent estimate of real mnemonic capacities, as it is a value obtained averaging four and not simply two values.

This study is not free from limitations. First, this is a cross-sectional study. Longitudinal investigations need to be planned to clarify the impact of the ApoE genotype along the axis of disease progression. Second, no specific control was carried out on other relevant risk factors involved in AD. Third, albeit these findings, surviving a Family-Wise Error-corrected cluster-level $p < 0.05$, show a degree of robustness against the Type I Error, larger samples are necessary to control further for more variables, and possibly, to investigate the impact of two copies of the ϵ_4 allele, since genotype effects might be attenuated in individuals carrying only one copy of the gene, and dose-dependent effects may exist. Fourth, no information on amyloid pathology was available for these patients. Although this would have not affected the diagnosis in these patients (as this had been reached after planned follow-up assessments over an extended period), it would have allowed us to understand whether the differences in connectivity shown by ϵ_4 carriers are associated with regional difference in amyloid deposition.

Despite these limitations, this study highlights the role of the ϵ_4 allele as a modifying-factor of neural pathways that are relevant for AD neurodegeneration. Although these findings shed some additional light on the role of the ϵ_4 allele in AD, the overall context in which these results have to be inserted remains extremely complicated. The clinical role of the hippocampus is not confined to the sole AD, but is involved in other conditions of neurological relevance. For example, alterations of resting-state hippocampal blood perfusion has been described in individuals at high risk of psychotic symptoms [70], and abnormal increases of hippocampal connectivity are visible as a consequence of post-traumatic brain disorders [71]. At the same time, the ϵ_4 allele was found to influence the activity and network properties of other, non-hippocampal regions, such as the prefrontal cortex [72] or midline structures and insula [73]. Finally, it is still unclear how the interplay of ApoE genotype and disease mechanisms influence not only resting-state by also task-based functional connectivity, especially during memory retrieval processes [51].

Conclusion

In summary, we found evidence of network spatial discrepancies between prodromal AD patients carrying one copy of the ApoE ϵ_4 allele and patients homozygous for the ApoE ϵ_3 allele showing no differences in demographic, neurostructural, and neuropsychological characteristics. Increased

connectivity was seen in carriers between the left hippocampus and 1) parietal areas, 2) prefrontal regions, and 3) the insular cortex. Although this pattern may be partially seen as the result of compensatory mechanisms (especially with regard to the increased connectivity seen in the frontal lobe), there are two interpretational avenues that identify these differences as maladaptive. First the “hyperfunctional hippocampus” hypothesis suggests that the excessive seed-to-parietal connectivity might be dysfunctional as negatively associated with cognitive performance [48-49]. This eventuality was ruled out by post-hoc analyses, which confirmed that the magnitude of connectivity along this pathway was positively associated with memory performance (characterising it as compensatory). Second, the “dishinhibition hypothesis” indicates that excessive hippocampus-to-insula connectivity might be result of maladaptive rewiring of a portion of the DMN with a portion of the salience network [50, 52]. As a consequence, the presence of maladaptive processes and the necessity to rely on extra compensation indicate that ϵ_4 carriers show an accentuated degree of network involvement independent of the diagnostic group and dependent on the ApoE genotype.

Conflict of Interest

The authors declare that they have no conflict of interest.

References

- [1] Andrews-Hanna JR, Reidler JS, Sepulcre J, Poulin R, Buckner RL. Functional-anatomic fractioning of the brain’s default network. *Neuron* 65(4): 550-562 (2010).
- [2] Buckner RL, Andrews-Hanna JR, Schacter DL. The brain’s default network. *Ann NY Acad Sci* 1124: 1-38 (2008).

- [3] Greicius MD, Srivastava G, Reiss AL, Menon V. Default-mode network activity distinguishes Alzheimer's disease from healthy aging: Evidence from functional MRI. *Proc Nat Acad Sci U S A* 101(13): 4637-4642 (2004).
- [4] Damoiseaux JS, Prater KE, Miller BL, Greicius MD. Functional connectivity tracks clinical deterioration in Alzheimer's disease. *Neurobiol Aging* 33: 828.e19-828.e30 (2012).
- [5] Jones DT, Machulda MM, Vemuri P, McDade EM, Zeng G, Senjem ML, et al. Age-related changes in the default mode network are more advanced in Alzheimer's disease. *Neurology* 77(16):1524-1531 (2011).
- [6] Koch W, Teipel S, Mueller S, Benninghoff J, Wagner M, Bokde ALW, et al. Diagnostic power of default mode network resting state fMRI in the detection of Alzheimer's disease. *Neurobiol Aging* 33: 466-478 (2012).
- [7] Qi Z, Wu X, Wang Z, Zhang N, Dong H, Yao L, et al. Impairment and compensation coexist in amnesic MCI default mode network. *Neuroimage* 50(1): 48-55 (2010).
- [8] Mueller SG, Schuff N, Yaffe K, Madison C, Miller B, Weiner MW. Hippocampal atrophy patterns in mild cognitive impairment and Alzheimer's disease. *Hum Brain Mapp* 31(9): 1339-47 (2010).
- [9] Ferreira LK, Diniz BS, Forlenza OV, Busatto GF, Zanetti MV. Neurostructural predictors of Alzheimer's disease: a meta-analysis of VBM studies. *Neurobiol Aging* 32(10): 1733-1741 (2011).
- [10] Wang Z, Liang P, Jia X, Qi Z, Yu L, Yang Y, et al. Baseline and longitudinal patterns of hippocampal connectivity in mild cognitive impairment: evidence from resting state fMRI. *J Neurol Sci* 309(1-2): 79-85 (2011).
- [11] Sperling RA, Dickerson BC, Pihlajamaki M, Vannini P, LaViolette PS, Vitolo OV, et al. Functional alterations in memory networks in early Alzheimer's disease. *Neuromolecular Med* 12(1): 27-43 (2010).

- [12] Putcha D, Brickhouse M, O'Keefe K, Sullivan C, Rentz D, Marshall G, et al. Hippocampal hyperactivation associated with cortical thinning in Alzheimer's disease signature regions in non-demented elderly adults. *J Neurosci* 31(48): 17680-17688 (2011).
- [13] Lambert JC, Ibrahim-Verbaas CA, Harold D, Naj AC, Sims R, Bellenquez C, et al. Meta-analysis of 74,046 individuals identifies 11 new susceptibility loci for Alzheimer's disease. *Nat Genet* 45: 1452-1458 (2013).
- [14] Strittmatter WJ, Roses AD. Apolipoprotein E and Alzheimer's disease. *Proc Nat Acad Sci U S A* 92: 4725-4727 (1995).
- [15] Sando SB, Melquist S, Cannon A, Hutton ML, Sletvold O, Saltvedt I, et al. APOE ϵ 4 lowers age at onset and is a high risk factor for Alzheimer's disease; A case control study from central Norway. *BMC Neurol* 8(9): 1-7 (2008).
- [16] Mahley RW, Rall Jr. SC. Apolipoprotein E: Far more than a lipid transport protein. *Ann Rev Genomics Hum Genet* 1: 507-537 (2000).
- [17] Mahley RW, Weisgraber KH, Huang Y. Apolipoprotein E4: A causative factor and therapeutic target in neuropathology, including Alzheimer's disease. *Proc Nat Acad Sci U S A* 103(15): 5644-5651 (2006).
- [18] Liu CC, Kanekiyo T, Xu H, Bu G. Apolipoprotein E and Alzheimer's disease: risk, mechanisms and therapy. *Nat Rev Neurol* 9(2): 106-118 (2013).
- [19] Mattay VS, Goldberg TE, Sambataro F, Weinberger DR. Neurobiology of cognitive aging: Insight from imaging genetics; *Biol Psychol* 79: 9-22 (2008).
- [20] Filippini N, MacIntosh BJ, Hough MJ, Goodwin GM, Frisoni GB, Smith SM, et al. Distinct patterns of brain activity in young carriers of the APOE- ϵ 4 allele. *Proc Nat Acad Sci U S A* 106(17): 7209-7214 (2009).

- [21] Machulda MM, Jones DT, Vemuri P, McDade E, Avula R, Przybelski S, et al. Effect of APOE $\epsilon 4$ status on intrinsic network connectivity in cognitively normal elderly. *Arch Neurol* 68(9): 1131-1136 (2011).
- [22] Sheline YI, Morris JC, Snyder AZ, Price JL, Yan Z, D'Angelo G, et al. APOE4 allele disrupts resting state fMRI connectivity in the absence of amyloid plaques or decreased CSF A β 42. *J Neurosci* 30(50): 17035-17040 (2010).
- [23] Westlye ET, Lundervold A, Rootwelt H, Lundervold AJ, Westlye LT. Increased hippocampal default mode synchronization during rest in middle-aged and elderly APOE $\epsilon 4$ carriers: relationships with memory performance. *J Neurosci* 31(21): 7775-7783 (2011).
- [24] Yang AC, Huang CC, Liu ME, Liou YJ, Hong CJ, Lo MT, et al. The APOE $\epsilon 4$ allele affects complexity and functional connectivity of resting brain activity in healthy adults. *Hum Brain Mapp* 35(7): 3238-3248 (2013).
- [25] Heise V, Filippini N, Trachtenberg AJ, Suri S, Ebmeier KP, Mackay CE. Apolipoprotein E genotype, gender and age modulate connectivity of the hippocampus in healthy adults. *Neuroimage* 98: 23-30 (2014).
- [26] Shu H, Shi YM, Chen G, Wang Z, Liu D, Yue CX, et al. Opposite neural trajectories of Apolipoprotein E $\epsilon 4$ and $\epsilon 2$ alleles with aging associated with different risks of Alzheimer's disease. *Cereb Cortex*. Epub ahead of print (2014). Available from <http://cercor.oxfordjournals.org/content/early/2014/10/21/cercor.bhu237.long>
- [27] Matura S, Prvulovic D, Butz M, Hartmann D, Sepanski B, Linnemann K, et al. Recognition memory is associated with altered resting-state functional connectivity in people at genetic risk for Alzheimer's disease. *Eur J Neurosci* 40(7): 3128-3135 (2014).
- [28] Song HQ, Long HX, Zuo XM, Yu CS, Liu B, Wang ZQ, et al. APOE Effects on Default Mode Network in Chinese Cognitive Normal Elderly: Relationship with Clinical Cognitive Performance. *PLoS One*. 10(7): e0133179 (2015).

- [29] Jelic C, Julin P, Shigeta M, Nordberg A, Lannfelt L, Winblad B, et al. Apolipoprotein E ϵ 4 allele decreases functional connectivity in Alzheimer's disease as measured by EEG coherence. *J Neurol Neurosurg Psychiatry* 63(1): 59-65 (1997).
- [30] Kramer G, van der Flier WM, de Langen C, Blankenstein MA, Scheltens P, Stam CJ. EEG functional connectivity and ApoE genotype in Alzheimer's disease and controls. *Clin Neurophysiol* 119: 2727-2732 (2008).
- [31] Canuet L, Tellado I, Couceiro V, Fraile C, Fernandez-Novoa L, Ishii R, et al. Resting-state network disruption and APOE genotype in Alzheimer's disease: A lagged functional connectivity study. *PLoS One* 7(9): e46289 (2013).
- [32] Cuesta P, Garcés P, Castellanos NP, López ME, Aurtenetxe S, Bajo R, et al. Influence of the APOE ϵ 4 allele and mild cognitive impairment diagnosis in the disruption of the MEG resting state functional connectivity in sources space. *J Alzheimers Dis* 44(2): 493-505 (2015).
- [33] Wang X, Wang JH, He Y, Li HY, Yuan HS, Evans A, et al. Apolipoprotein E ϵ 4 modulates cognitive profiles, hippocampal volume, and resting-state functional connectivity in Alzheimer's disease. *J Alzheimers Dis* 45(3): 781-795 (2015).
- [34] Wang JH, Wang X, He Y, Yu X, Wang HL, He Y. Apolipoprotein E ϵ 4 modulates functional brain connectome in Alzheimer's disease. *Hum Brain Mapp* 36(5): 1828-1846 (2015).
- [35] Belleville S, Clement F, Mellah S, Gilbert B, Fontaine F, Gauthier S. Training-related brain plasticity in subjects at risk of developing Alzheimer's disease. *Brain* 134: 1623-1634 (2011).
- [36] Rosen AC, Sugiura L, Kramer JH, Whitfield-Gabrieli S, Gabrieli JD. Cognitive training changes hippocampal function in mild cognitive impairment: A pilot study. *J Alzheimers Dis* 26: 349-357 (2011).
- [37] Dubois B, Feldman HH, Jacova C, Dekosky ST, Barberger-Gateau P, Cummings J, et al. Research criteria for the diagnosis of Alzheimer's disease: revising the NINCDS-ADRDA criteria. *Lancet Neurol* 6(8): 734-746 (2006).

- [38] Petersen RC, Doody R, Kurz A, Mohs RC, Morris JC, Rabins PV, et al. Current concepts in mild cognitive impairment. *Arch Neurol* 58(12): 1985-1992 (2001).
- [39] Ashburner J, Friston KJ. Voxel-based morphometry--the methods. *Neuroimage* 11(6): 805-821 (2000).
- [40] Jorge Cardoso M, Leung K, Modat M, Keihaninejad S, Cash D, Barnes J, et al. STEPS: Similarity and Truth Estimation for Propagated Segmentations and its application to hippocampal segmentation and brain parcellation. *Med Image Anal* 17(6): 671-684 (2013).
- [41] Song XW, Dong ZY, Long XY, Li SF, Zuo XN, Zhu CZ, et al. REST: A toolkit for resting-state functional magnetic resonance imaging data processing. *PLoS One* 6(9): e25031 (2011).
- [42] Maldjian JA, Laurienti PJ, Kraft RA, Burdette JH. An automated method for neuroanatomic and cytoarchitectonic atlas-based interrogation of fmri data sets. *Neuroimage* 19: 1233–1239 (2003).
- [43] Brett M, Anton JL, Valabregue R, Poline JB. Region of interest analysis using an SPM toolbox. Available on CD-ROM in *Neuroimage* 16(2) (2002).
- [44] Wang L, Zang Y, He Y, Liang M, Zhang X, Tian L, et al. Changes in hippocampal connectivity in the early stages of Alzheimer's disease: evidence from resting state fMRI. *Neuroimage* 31(2): 496-504 (2006).
- [45] Allen G, Barnard H, McColl R, Hester AL, Fields JA, Weiner MF, et al. Reduced hippocampal functional connectivity in Alzheimer disease. *Arch Neurol* 64(10): 1482-1487 (2007).
- [46] Lancaster JL, Summerlin JL, Rainey L, Freitas CS, Fox PT. The Talairach Daemon, a database server for Talairach Atlas Labels. *Neuroimage* 5: S633 (1997).
- [47] Lancaster JL, Woldorff MG, Parsons LM, Liotti M, Freitas CS, Rainey L, et al. Automated Talairach atlas labels for functional brain mapping. *Hum Brain Mapp* 10: 120–131 (2000).

- [48] Tahmasian M, Pasquini L, Scherr M, Meng C, Förster S, Mulej Bratec S, et al. The lower hippocampus global connectivity, the higher its local metabolism in Alzheimer disease. *Neurology* 84(19): 1956-1963 (2015).
- [49] Trachtenberg AJ, Filippini N, Mackay CE. The effects of APOE- ϵ 4 on the BOLD response. *Neurobiol Aging* 33: 323-334 (2012).
- [50] Bakker A, Krauss GL, Albert MS, Speck CL, Jones LR, Stark CE, et al. Reduction of hippocampal hyperactivity improves cognition in amnesic mild cognitive impairment. *Neuron* 10;74(3): 467-474 (2012).
- [51] Bai F, Zhang ZJ, Watson DR, Yu H, Shi YM, Yuan YG, et al. Abnormal functional connectivity of hippocampus during episodic memory retrieval processing network in amnesic mild cognitive impairment. *Biol Psychiatry* 65(11): 951-958 (2009).
- [52] Seeley WW. Divergent network connectivity changes in healthy ApoE ϵ 4 carriers. *Arch Neurol* 68(9): 1107-1108 (2011).
- [53] Seeley WW, Menon V, Schatzberg AF, Keller J, Glover GH, Kenna H, et al. Dissociable intrinsic connectivity networks for salience processing and executive control. *J Neurosci* 27(9): 2349-2356 (2007).
- [54] Goveas JS, Xie CM, Chen G, Li WJ, Ward BD, Franczak MB. Functional network endophenotypes unravel the effects of apolipoprotein E epsilon 4 in middle-aged adults. *PLoS One* 8(2): e55902 (2013).
- [55] Gardini S, Cueto F, Fasano F, Ferrari Pellegrini F, Marchi M, Venneri A, et al. Brain structural substrates of semantic memory decline in mild cognitive impairment. *Curr Alzheimer Res* 10(4): 373-389 (2013).
- [56] Rodríguez-Ferreiro J, Cueto F, Monsalve A, Martínez C, Pérez AJ, Venneri A. Establishing the relationship between cortical atrophy and semantic deficits in Alzheimer's disease and mild cognitive impairment patients through voxel-based morphometry. *J Neurolinguistics* 25: 139-149 (2012).

- [57] Dem Heijer T, Oudkerk M, Launer LJ, van Duijn CM, Hofman A, Breteler MMB. Hippocampal, amygdalar, and global brain atrophy in different apolipoprotein E genotypes. *Neurology* 59(5): 746-748 (2002).
- [58] Farlow MR, He Y, Tekin S, Xu J, Lane R, Charles HC. Impact of APOE in mild cognitive impairment, *Neurology* 63(10): 1898-1901 (2004).
- [59] Fleisher A, Grundman M, Jack CR Jr, Petersen RC, Taylor C, Kim HT, et al. Sex, apolipoprotein E epsilon status, and hippocampal volume in mild cognitive impairment. *Arch Neurol* 62(6): 953-957 (2005).
- [60] Pennanen C, Testa C, Boccardi M, Laakso MP, Hallikainen M, Helkala EL, et al. The effect of apolipoprotein polymorphism on brain in mild cognitive impairment: a voxel-based morphometric study. *Dem Geriatr Cogn Disord* 22(1): 60-66 (2006).
- [61] Liu Y, Pajanen T, Westman E, Wahlund LO, Simmons A, Tunnard C, et al. Effect of APOE ϵ 4 allele on cortical thicknesses and volumes: The AddNeuroMed study. *J Alzheimers Dis* 21: 947-966 (2010).
- [62] Spampinato MV, Rumboldt Z, Hosker RJ, Mintzer JE; Alzheimer's Disease Neuroimaging Initiative. Apolipoprotein E and gray matter volume loss in patients with mild cognitive impairment and Alzheimer disease. *Radiology* 258(3): 843-852 (2011).
- [63] Thomann PA, Roth AS, Dos Santos V, Toro P, Essig M, Schröder J. Apolipoprotein E polymorphism and brain morphology in mild cognitive impairment. *Dem Geriatr Cogn Disord* 26(4): 300-305 (2008).
- [64] Venneri A, McGeown WJ, Biundo R, Mion M, Nichelli P, Shanks MF. The neuroanatomical substrate of lexical-semantic decline in MCI APOE epsilon 4 carriers and noncarriers. *Alz Dis Assoc Dis* 25(3): 230-241 (2011).

- [65] Mosconi L, Perani D, Sorbi S, Herholz K, Nacmias B, Holthoff V, et al. MCI conversion to dementia and the APOE genotype: a prediction study with FDG-PET. *Neurology* 63(12): 2332-2340 (2004).
- [66] Pievani M, de Haan W, Wu T, Seeley WW, Frisoni GB. Functional network disruption in the degenerative dementias. *Lancet Neurol* 10: 829-843 (2011).
- [67] Kljajevic V, Meyer P, Holzmann C, Dyrba M, Kasper E, Bokde AL, et al. The $\epsilon 4$ genotype of apolipoprotein E and white matter integrity in Alzheimer's disease. *Alzheimers Dement* 10(3): 401-404 (2014).
- [68] Wang CS, Stebbins GT, Mediana DA, Shah RC, Bammer R, Moseley ME. et al. Atrophy and dysfunction of parahippocampal white matter in mild Alzheimer's disease. *Neurobiol Aging* 33(1): 43-52 (2012).
- [69] Bonner-Jackson A, Mahmoud S, Miller J, Banks SJ. Verbal and non-verbal memory and hippocampal volumes in a memory clinic population. *Alzheimers Res Ther* 7(1): 61 (2015).
- [70] Allen P, Chaddock CA, Egerton A, Howes OD, Bonoldi I, Zelaya F, et al. Resting hyperperfusion of the hippocampus, midbrain, and basal ganglia in people at high risk for psychosis. *Am J Psychiatry* 173(4): 392-399 (2016).
- [71] Dunkley BT, Doesburg SM, Sedge PA, Grodecki RJ, Shek PN, Pang EW, et al. Resting-state hippocampal connectivity correlates with symptom severity in post-traumatic stress disorder. *Neuroimage Clin* 5: 377-384 (2014).
- [72] De Marco M, Clough PJ, Dyer CE, Vince RV, Waby JS, Midgley AW, et al. Apolipoprotein E $\epsilon 4$ allele modulates the immediate impact of acute exercise on prefrontal function. *Behav Genet* 45(1):106-116 (2015).
- [73] Yao Z, Hu B, Zheng J, Zheng W, Chen X, Gao X, et al. A FDG-PET Study of Metabolic Networks in Apolipoprotein E $\epsilon 4$ Allele Carriers. *PLoS One* 10(7): e0132300 (2015).

Figure Captions

Fig.1

Pattern of enhanced functional connectivity of the left hippocampus found in patients who were carriers of the ϵ_4 allele. Slices in MNI space are as follows: $z = -10$, $z = 46$. The seed region is illustrated on the left (slices in MNI space are as follows: $x = -26$, $y = -13$, $z = -13$).

Fig.2

Association between mediotemporal-parietal connectivity and memory function as investigated at post-hoc. On the left, the association found in the group of patients who were homozygotes for the ϵ_3 allele ($r = -0.037$; $p = 0.905$); on the right, the association found in the group of patients with an $\epsilon_4\epsilon_3$ genotype ($r = 0.602$; $p = 0.029$).




Figure 1




Figure 2

Table 1: Sample Characterisation

Descriptive Variables	£4£3	£3£3	Between-Group Statistics	
Demographic Characteristics			Pt test/chi square	
Age at Scan (years)	73.20 (5.85)	73.80 (4.87)	0.762	
Education Level (years)	9.20 (4.11)	9.13 (3.56)	0.962	
Gender (f/m)	7/8	7/8	0.999	
Global Structural Indices			Pt test	
Grey-Matter Volume (cl)	538.74 (64.45)	536.66 (59.93)	0.893	
White-Matter Volume (cl)	451.09 (65.14)	423.83 (63.69)	0.256	
Brain Parenchymal Volume (cl)	990.83 (123.53)	960.49 (114.58)	0.491	
Cerebrospinal-Fluid Volume (cl)	697.05 (122.87)	770.58 (112.12)	0.098	
Grey-Matter Fraction	0.32 (0.02)	0.31 (0.03)	0.244	
White-Matter Fraction	0.27 (0.02)	0.24 (0.02)	0.002	
Brain Parenchymal Fraction	0.59 (0.03)	0.56 (0.04)	0.012	
Total Intracranial Volume (cl)	1687.88 (231.15)	1731.07 (181.37)	0.574	
Left Hippocampal Volume (cl)	2.12 (0.37)	2.24 (0.27)	0.322	
Right Hippocampal Volume (cl)	2.24 (0.30)	2.30 (0.30)	0.543	
Left Hippocampal Fraction	2.15 ⁻³ (2.95 ⁻⁴)	2.35 ⁻³ (2.85 ⁻⁴)	0.065	
Right Hippocampal Fraction	2.27 ⁻³ (2.82 ⁻⁴)	2.41 ⁻³ (2.99 ⁻⁴)	0.190	
Hippocampal Asymmetry (left/right)	0.95 (0.10)	0.97 (0.04)	0.356	
Neuropsychological Raw Scores			Pt test	P_{Corrected ANOVA}
Mini Mental-State Examination	25.93 (3.81)	26.00 (3.21)	0.959	0.921
Raven Progressive Matrices	26.33 (5.43)	25.00 (6.47)	0.546	0.501
Letter Fluency Test	33.00 (11.93)	27.60 (10.36)	0.196	0.163
Category Fluency Test	27.07 (8.52)	25.47 (6.62)	0.570	0.645
Digit Cancellation Test	48.00 (9.18)	46.67 (9.40)	0.697	0.729
WAIS - Similarities	17.93 (5.59)	17.13 (5.29)	0.690	0.686
Token Test	33.57 (2.00)	33.50 (2.10)	0.930	0.927
Rey Complex Figure Test - Copy	29.23 (3.95)	28.13 (6.36)	0.574	0.601
Rey Complex Figure Test - Recall	8.40 (4.40)	7.87 (3.92)	0.728	0.809
Stroop Time Interference Effect	36.43 (20.32)	40.43 (14.04)	0.536	0.547
Stroop Error Interference Effect	1.83 (2.74)	2.60 (2.83)	0.457	0.515
Digit Span Test - Forward	5.40 (0.74)	5.80 (0.86)	0.183	0.199
Digit Span Test - Backwards	3.80 (0.86)	3.67 (0.62)	0.630	0.547
Corsi Test	4.13 (0.74)	4.13 (0.74)	0.999	0.876
Visual Supraspan Test	14.91 (7.62)	9.83 (7.66)	0.085	0.091
Prose Memory Test - Immediate Recall	5.67 (3.73)	5.93 (3.15)	0.834	0.731
Prose Memory Test - Delayed Recall	6.20 (4.84)	6.53 (3.85)	0.836	0.772
Prose Memory Test - Global Recall	11.87 (8.17)	12.47 (6.60)	0.826	0.739
Paired Associates Test	9.61 (4.27)	7.60 (2.68)	0.139	0.159
Confrontation Naming Test	18.07 (2.12)	18.00 (1.65)	0.924	0.972

Neuropsychological Corrected Scores			p_{t test}	Cut-Off
Raven Progressive Matrices	28.71 (4.28)	28.05 (6.09)	0.736	≤ 18.96
Letter Fluency Test	37.47 (9.81)	31.87 (11.33)	0.159	≤ 16
Category Fluency Test	32.00 (9.14)	31.07 (6.70)	0.752	≤ 24
Digit Cancellation Test	49.22 (7.18)	46.32 (9.25)	0.346	≤ 30
Token Test	31.65 (7.33)	33.68 (1.99)	0.309	≤ 26.25
Rey Complex Figure Test - Copy	30.85 (3.71)	29.23 (6.64)	0.419	≤ 28.87
Rey Complex Figure Test - Recall	11.93 (4.85)	12.20 (4.18)	0.873	≤ 9.46
Stroop Time Interference Effect	26.38 (19.43)	29.48 (14.03)	0.620	≥ 36.92
Stroop Error Interference Effect	1.02 (2.25)	1.52 (2.21)	0.544	≥ 4.24
Digit Span Test - Forward	5.57 (0.69)	6.02 (0.96)	0.151	≤ 3.5
Corsi Test	4.42 (0.74)	4.50 (0.63)	0.742	≤ 3.25
Visual Supraspan Test	16.89 (7.24)	11.48 (7.77)	0.063	≤ 5.5
Prose Memory Test - Global Recall	<u>12.67 (7.29)</u>	<u>13.07 (6.69)</u>	0.877	≤ 15.76
Paired Associates Test	11.14 (3.81)	9.37 (3.01)	0.174	≤ 6

Between-group statistics were run using chi square (gender), one-way ANOVA (differences in cognitive performance corrected for age and levels of education) and independent-sample t (all remaining comparisons) inferential models. Hippocampal asymmetry was calculated based on the raw volumes computed using the STEPS protocol. Brain parenchymal volume was computed as the sum of grey-matter and white-matter volumes. Tissue fractions were calculated dividing tissue class volume by total intracranial volume. Hippocampal fractions were instead calculated dividing hippocampal volumes by brain parenchymal volumes. “pCorrected ANOVA” indicates the significance level after covarying for years of age and years of education. In bold the sole between-group difference surviving Bonferroni correction for multiple comparisons (pGlobal Structural Indices < 0.0038). The aspects of cognitive functions showing performance below cut-off are instead underlined.

Table 2: Between-group differences in hippocampal connectivity

Cluster Number	Cluster Size (voxels)	Cluster-Level pFWE	Z Value at Local Maximum	Hemisphere	Cerebral Region	Brodmann Area	Talairach Coordinates		
							x	y	z
1	377	0.047	3.72	L	Inferior Frontal Gyrus	47	-38	15	-9
			3.44	L	Insula	13	-46	12	-1
			2.62	L	Inferior Frontal Gyrus	13	-32	13	-14
2	506	0.010	3.25	R	Inferior Parietal Lobule	40	40	-43	43
			3.21	R	Inferior Parietal Lobule	40	38	-52	45
			3.18	R	Postcentral Gyrus	2	59	-29	42

Table 3: Characteristics of the sample of patients extracted from the ADNI cohort

Descriptive Variables	ε4ε3 - ε4ε4	ε3ε3	Between-Group Statistics
Age at Recruitment (years)	71.43 (5.61)	72.43 (7.69)	0.559
Education Level (years)	16.70 (2.88)	15.97 (2.42)	0.272
Gender (f/m)	12/18	17/18	0.488
Ventricular Size (mm ³)	36785.23 (20974.18)	37645.34 (240505.53)	0.879
Mini Mental State Examination	27.50 (1.96)	28.00 (1.61)	0.263
Rey Auditory Verbal Learning Test - Immediate Recall	36.53 (9.53)	34.34 (9.90)	0.604
Rey Auditory Verbal Learning Test - Learning	4.37 (3.18)	4.89 (2.98)	0.318

Between-group statistics were run using chi square (gender), one-way ANOVA (differences in cognitive performance corrected for age and levels of education) and independent-sample t (all remaining comparisons)
inferential models